VOL. 749, NOS. 1+2 JOURNAL OF CHROMATOGRAPHY A 18 OCTOBER 1996

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

OBITUARIES

Emanuel Gil-Av
by N. Grinberg (Rahway, NJ, USA) 1

K.I. Sakodynskii
by V.A. Davankov (Moscow, Russia) 3

REGULAR PAPERS

General

Artificial neural networks aided deconvolving overlapped peaks in chromatograms
by H. Miao, M. Yu and S. Hu (Hangzhou, China) (Received 26 March 1996) 5

Column Liquid Chromatography

Immobilized liposome chromatography of drugs on capillary continuous beds for model analysis of drug–membrane interactions
by Y. Zhang, C.-M. Zeng, Y.-M. Li, S. Hjertén and P. Lundahl (Uppsala, Sweden) (Received 29 April 1996) 13

On the relevance of face-to-edge $\pi-\pi$ interactions to chiral recognition
by W.H. Pirkle and Y. Liu (Urbana, IL, USA) (Received 7 May 1996) 19

Enantioseparation in the synthesis of *myo*-inositol phosphates by high-performance liquid chromatography using a β-cyclodextrin-bonded column
by A. Burmester and B. Jastorff (Bremen, Germany) (Received 9 April 1996) 25

Isolation of dinophysistoxin-2 and the high-performance liquid chromatographic analysis of diarrhetic shellfish toxins using derivatisation with 1-bromoacetylpyrene
by S.S. Kelly, A.G. Bishop, E.P. Carmody and K.J. James (Cork, Ireland) (Received 26 February 1996) 33

Rapid method for the determination of the substitution pattern of O-methylated 1,4-glucans by high-pH anion-exchange chromatography with pulsed amperometric detection
by J. Heinrich and P. Mischnick (Hamburg, Germany) (Received 7 May 1996) 41

Mechanistic aspects of fatty acid retention in silver ion chromatography
by B. Nikolova-Damyanova and W.W. Christie (Dundee, UK) and B. Herslöf (Stockholm, Sweden) (Received 11 April 1996) 47

Rapid, quantitative determination of polar compounds in fats and oils by solid-phase extraction and size-exclusion chromatography using monostearin as internal standard
by G. Márquez-Ruiz, N. Jorge, M. Martín-Polvillo and M.C. Dobarganes (Sevilla, Spain) (Received 24 April 1996) . 55

Resolution of enantiomeric steroids by high-performance liquid chromatography on chiral stationary phases
by M. Kummer and H.-J. Palme (Jena, Germany) and G. Werner (Leipzig, Germany) (Received 4 April 1996) . . . 61

Thiol pegylation facilitates purification of chymopapain leading to diffraction studies at 1.4 Å resolution
by M. Azarkan (Brussels, Belgium), D. Maes, J. Bouckaert, M.-H.D. Thi and L. Wyns (Sint Genesius Rode, Belgium) and Y. Looze (Brussels, Belgium) (Received 10 April 1996) 69

Integrated derivatization–chemiluminescence detection system for the determination of β-carboline alkaloids by high-performance liquid chromatography
by J. Cepas, M. Silva and D. Pérez-Bendito (Córdoba, Spain) (Received 4 April 1996) 73

Liquid chromatographic method for the determination of nicotine in pharmaceutical formulations
by A.K. Dash and S.-T. Wong (Omaha, NE, USA) (Received 10 April 1996) 81

(Continued overleaf)

Contents (continued)

Separation of zinc dialkyldithiophosphates in lubricating oil additives by normal-phase high-performance liquid chromatography
by N. Lambropoulos, T.J. Cardwell and D. Caridi (Bundoora, Australia) and P.J. Marriott (Melbourne, Australia) (Received 26 March 1996) 87

High-performance liquid chromatographic determination of 2-furaldehyde and 5-hydroxymethyl-2-furaldehyde in honey
by F.L. Coco (Udine, Italy), C. Valentini (Mestre, Italy), V. Novelli (Udine, Italy) and L. Ceccon (Trieste, Italy) (Received 19 April 1996) 95

Determination of trace levels of thorium(IV) and uranyl by reversed-phase chromatography with on-line preconcentration and ligand exchange
by F. Hao, B. Paull and P. R. Haddad (Hobart, Australia) (Received 16 April 1996) 103

Ion chromatographic separation of selenite and selenate using a polyanionic eluent
by P. Janoš and P. Aczel (Brno, Czech Republic) (Received 23 April 1996) 115

Gas Chromatography

New method for estimating vapor pressure by the use of gas chromatography
by S.F. Donovan (Princeton, NJ, USA) (Received 23 April 1996) 123

Relative responses of various classes of compounds using a pulsed discharge helium photoionization detector. Experimental determination and theoretical calculations
by S. Mendonca, W.E. Wentworth, E.C.M. Chen and S.D. Stearns (Houston, TX, USA) (Received 26 March 1996) . 131

Experimental and theoretical relative response factors for a pulsed discharge krypton photoionization detector
by W.E. Wentworth, S. Watanesk, N. Helias, R. Swatloski, E.C.M. Chen and S.D. Stearns (Houston, TX, USA) (Received 26 March 1996) 149

Optimization of solid-phase extraction method for analysis of low-ppb amounts of aldehydes-ozonation by-products
by J. Nawrocki, I. Kalkowska and A. Dąbrowska (Poznań, Poland) (Received 9 April 1996) 157

Simultaneous determination of chlorophenols, chlorobenzenes and chlorobenzoates in microbial solutions using pentafluorobenzylbromide derivatization and analysis by gas chromatography with electron-capture detection
by C.L. Gabelish, P. Crisp and R.P. Schneider (Sydney, Australia) (Received 24 April 1996) 165

Lesser alkaloids of cocaine-bearing plants. II. 3-Oxo-substituted tropane esters: detection and mass spectral characterization of minor alkaloids found in South American *Erythroxylum coca* var. *coca*
by J.F. Casale and J.M. Moore (McLean, VA, USA) (Received 6 May 1996) 173

Screening method for the determination of 28 volatile organic compounds in indoor and outdoor air at environmental concentrations using dual-column capillary gas chromatography with tandem electron-capture–flame ionization detection
by J. Begerow, E. Jermann, T. Keles, T. Koch and L. Dunemann (Düsseldorf, Germany) (Received 23 April 1996) . 181

Supercritical Fluid Chromatography

Packed-column supercritical fluid chromatography of a new dihydropyridine drug based on direct injection of emulsion samples
by L. Karlsson, O. Gyllenhaal, A. Karlsson and J. Gottfries (Mölndal, Sweden) (Received 9 April 1996) 193

Planar Chromatography

Solute effects on reversed-phase thin-layer chromatography. A linear free energy relationship analysis
by M.H. Abraham, C.F. Poole and S.K. Poole (London, UK) (Received 16 April 1996) 201

Novel computational methods for the determination of partition coefficients by planar chromatography
by R.M. Kleyle, D. Nurok, A.D. Kossoy and S.C. Burris (Indianapolis, IN, USA) (Received 24 April 1996) 211

Electrophoresis

On-line partial filling micellar electrokinetic chromatography–electrospray ionization mass spectrometry
by W.M. Nelson, Q. Tang, A.K. Harrata and C.S. Lee (Ames, IA, USA) (Received 29 April 1996) 219

Separation of structural homologues of alkylphenols and isomers of 4-nonylphenol by cyclodextrin-modified micellar electrokinetic chromatography
by Y. He and H. K. Lee (Kent Ridge, Singapore) (Received 26 March 1996) 227

Contents (continued)

Capillary electrophoretic examination of underivatized oligosaccharide mixtures released from immunoglobulin G antibodies and CTLA4Ig fusion protein
by K.F. Greve and D.E. Hughes (Syracuse, NY, USA) and B.L. Karger (Boston, MA, USA) (Received 2 April 1996) 237

Separation of pharmaceutically important estrogens by micellar electrokinetic chromatography
by S.K. Poole (London, UK) and C.F. Poole (London, UK and Detroit, MI, USA) (Received 17 April 1996) 247

Utilization of *n*-alkyl-β-D-glucopyranosides in enantiomeric separation by micellar electrokinetic chromatography
by P.L. Desbène and C.E. Fulchic (Evreux and Mont Saint Aignan, France) (Received 19 April 1996) 257

Separation of toxic peptides (microcystins) in capillary electrophoresis, with the aid of organic mobile phase modifiers
by N. Onyewuenyi and P. Hawkins (West Ryde, Australia) (Received 9 April 1996) 271

Control of separation selectivity and electroosmotic flow in nonaqueous capillary electrophoretic separations of alkali and alkaline earth metal ions
by H. Salimi-Moosavi and R.M. Cassidy (Saskatoon, Canada) (Received 28 February 1996) 279

More sensitive way to determine iron using an iron(II)–1,10-phenanthroline complex and capillary electrophoresis
by J. Xu, P. Che and Y. Ma (Kirksville, MO, USA) (Received 19 April 1996) 287

SHORT COMMUNICATIONS

Column Liquid Chromatography

Simultaneous analysis of individual catechins and caffeine in green tea
by T. Goto (Ibaraki-ken, Japan), Y. Yoshida and M. Kiso (Tokyo, Japan) and H. Nagashima (Ibaraki-ken, Japan) (Received 29 April 1996) 295

Electrophoresis

Cyclodextrins as buffer additives for the enantiomeric separation of pinacidil in capillary zone electrophoresis
by Z. Wang, A. Huang, Y. Sun and Z. Sun (Beijing, P.R. China) (Received 25 April 1996) 300

AUTHOR INDEX VOL. 749 304